

CYCLE #2 Simplifying Expressions and Algebraic Gymnastics

Show work / No Calculator

1. Find the exact value of each expression

(a) $\log_{10} 25 + \log_{10} 4$

(b) $e^{4\ln 2}$

2. Solve each of the following equations for x . Find the simplified, exact value.

(a) $e^x = 3$

(b) $e^{e^x} = 3$


(c) $\log_3(x+1) = 2$

(d) $\log_3 27 = x$

Multiple Choice

- _____ 3. Rationalize the numerator of $\frac{\sqrt{x+4} - \sqrt{x-2}}{x}$
- (A) $\frac{2}{x(\sqrt{x+4} + \sqrt{x-2})}$ (B) $\frac{6}{x(\sqrt{x+4} - \sqrt{x-2})}$ (C) $\frac{6}{x(\sqrt{x+4} + \sqrt{x-2})}$
(D) $\frac{2x}{\sqrt{x+4} + \sqrt{x-2}}$ (E) $\frac{6x}{\sqrt{x+4} - \sqrt{x-2}}$

- ____ 4. Which, if any, of the following statements are true when a, b are real numbers?
- For all positive a and b , $\sqrt{a+b} = \sqrt{a} + \sqrt{b}$.
 - For all a and b , $\sqrt{(a+b)^2} = |a+b|$.
 - For all positive a and b , $\frac{a-b}{\sqrt{a} + \sqrt{b}} = \sqrt{a} - \sqrt{b}$.
- (A) III only (B) all of them (C) I and II only (D) II only (E) II and III only
 (F) none of them (G) I and III only (H) I only
- ____ 5. Simplify the expression $\frac{1+\frac{2}{x-3}}{5+40\left(\frac{x}{x^2-9}\right)}$
- (A) $\frac{1}{5}\left(\frac{x+3}{2x+9}\right)$ (B) $\frac{x+3}{x-9}$ (C) $\frac{1}{5}\left(\frac{x+3}{x+9}\right)$ (D) $\frac{x+3}{2x-9}$ (E) $\frac{1}{5}\left(\frac{x-3}{x+9}\right)$ (F) $\frac{x-3}{x-9}$
- ____ 6. The shaded area in the figure is the complement of the sector of a circle of radius 6 inches lying inside the right triangle ΔABC with the angle θ being expressed in radians. Express this shaded area as a function of S , of θ .




- (A) $S(\theta) = 36(\tan \theta - \theta)$ (B) $S(\theta) = 36(\sin \theta - \theta)$ (C) $S(\theta) = 18(\sin \theta - \theta)$
 (D) $S(\theta) = 18(\cos \theta - \theta)$ (E) $S(\theta) = 18(\tan \theta - \theta)$

_____ 7. Which of the following statements are true?

- I. The circle $(x-1)^2 + (y-2)^2 = 1$ has radius = 1.
 - II. The circle $(x-5)^2 + (y-6)^2 = 9$ has center = $(6, 5)$.
 - III. The circle $(x-4)^2 + (y-4)^2 = 25$ has y-intercepts = 1, 7.
- (A) I only (B) II only (C) I and III only (D) III only (E) II and III only
(F) none of them (G) all of them (H) I and II only

_____ 8. Find the area of the shaded region shown outside the square and inside the circle when the area of the circle is 25π sq. units.



- (A) $5(4-\pi)$ sq. units (B) $5(\pi-1)$ sq. units (C) $25(\pi-2)$ sq. units
(D) $5(\pi-2)$ sq. units (E) $25(\pi-1)$ sq. units (F) $25(4-\pi)$ sq. units

- _____ 9. Simplify the difference quotient $\frac{f(x+h)-f(x)}{h}$, ($h \neq 0$), when $f(x)=2x^2-4x-4$.
(A) $4x+4+2h$ (B) $4x-4+2h$ (C) $2x+4+2h$ (D) $2x-4+2h$ (E) $4x-4$

- _____ 10. Captain Calculus can leap over tall buildings. When he does so, his height s (in feet) off the ground after t seconds is given by $s(t)=-t^2+7t+34$. For how many seconds is Captain Calculus more than 40 feet off the ground?

(A) 6 sec (B) $\frac{9}{2}$ sec (C) $\frac{11}{2}$ sec (D) $\frac{5}{2}$ sec (E) 5 sec

- _____ 11. If $f(x)=2x-1$ and $g(x)=x+3$, which of the following gives $(f \circ g)(2)$?
(A) 2 (B) 6 (C) 7 (D) 9 (E) 10

- _____ 12. Which of the following is a solution of the equation $2-3^x=-1$?
(A) $x=-2$ (B) $x=-1$ (C) $x=0$ (D) $x=1$ (E) No solution

____ 13. The length L of a rectangle is twice as long as its width W . Which of the following gives the area A of the rectangle as a function of its width?

- (A) $A(W) = 3W$ (B) $A(W) = \frac{1}{2}W^2$ (C) $A(W) = 2W^2$
(D) $A(W) = W^2 + 2W$ (E) $A(W) = W^2 - 2W$

____ 14. If $p(x) = (x+2)(x+k)$ and if the remainder is 12 when $p(x)$ is divided by $x-1$, then $k =$
(A) 2 (B) 3 (C) 6 (D) 11 (E) 13

____ 15. The set of all points (e^t, t) , where t is a real number, is the graph of $y =$

- (A) $\frac{1}{e^x}$ (B) $e^{1/x}$ (C) $xe^{1/x}$ (D) $\frac{1}{\ln x}$ (E) $\ln x$

____ 16. If $f(x) = \frac{4}{x-1}$ and $g(x) = 2x$, then the solutions of $f(g(x)) = g(f(x))$ is

- (A) $\left\{\frac{1}{3}\right\}$ (B) $\{2\}$ (C) $\{3\}$ (D) $\{-1, 2\}$ (E) $\left\{\frac{1}{3}, 2\right\}$

_____ 17. If the function f is defined by $f(x) = x^5 - 1$, then f^{-1} , the inverse function of f , is defined by

$$f^{-1}(x) =$$

- (A) $\frac{1}{\sqrt[5]{x+1}}$ (B) $\frac{1}{\sqrt[5]{x-1}}$ (C) $\sqrt[5]{x-1}$ (D) $\sqrt[5]{x} - 1$ (E) $\sqrt[5]{x+1}$

_____ 18. If a, b, c, d , and e are real numbers and $a \neq 0$, then the polynomial equation

$$ax^7 + bx^5 + cx^3 + dx + e = 0$$
 has

- (A) only one real root (B) at least one real root (C) an odd number of nonreal roots
(D) no real roots (E) no positive real roots

_____ 19. What are all values of k for which the graph of $y = x^3 - 3x^2 + k$ will have three distinct x -intercepts?

- (A) All $k > 0$ (B) All $k < 4$ (C) $k = 0, 4$ (D) $0 < k < 4$ (E) All k

_____ 20. If $f(g(x)) = x^3 + 3x^2 + 4x + 5$ and $g(x) = 5$, then $g(f(x)) =$

- (A) $5x^2 + 15x + 25$ (B) $5x^3 + 15x^2 + 20x + 25$ (C) 1125 (D) 225 (E) 5

- _____ 21. If $f(x) = 2x^3 + Ax^2 + Bx - 5$ and if $f(2) = 3$ and $f(-2) = -37$, what is the value of $A + B$?
(A) -6 (B) -3 (C) -1 (D) 2 (E) It cannot be determined from the information given

- _____ 22. Suppose that f is a function that is defined for all real numbers. Which of the following conditions assures that f has an inverse function?

- (A) The function f is periodic (B) The function f is symmetric with respect to the y -axis
(C) The function f is concave up (D) The function f is a strictly increasing function
The function f is continuous

- _____ 23. If $\log_a(2^a) = \frac{a}{4}$, then $a =$
(A) 2 (B) 4 (C) 8 (D) 16 (E) 32

- _____ 24. If $f(g(x)) = \ln(x^2 + 4)$, $f(x) = \ln(x^2)$, and $g(x) > 0$ for all real x , then $g(x) =$
(A) $\frac{1}{\sqrt{x^2 + 4}}$ (B) $\frac{1}{x^2 + 4}$ (C) $\sqrt{x^2 + 4}$ (D) $x^2 + 4$ (E) $x + 2$

_____ 25. If $\ln x - \ln\left(\frac{1}{x}\right) = 2$, then $x =$

- (A) $\frac{1}{e^2}$ (B) $\frac{1}{e}$ (C) e (D) $2e$ (E) e^2

_____ 26. If $f(x) = \frac{x}{x+1}$, then the inverse function, f^{-1} , is given by $f^{-1}(x) =$

- (A) $\frac{x-1}{x}$ (B) $\frac{x+1}{x}$ (C) $\frac{x}{1-x}$ (D) $\frac{x}{x+1}$ (E) x

_____ 27. If $f(x) = e^x \sin x$, then the number of zeros of f on the closed interval $[0, 2\pi]$ is

- (A) 0 (B) 1 (C) 2 (D) 3 (E) 4

_____ 28. If h is the function given by $h(x) = f(g(x))$, where $f(x) = 3x^2 - 1$ and $g(x) = |x|$, then
 $h(x) =$

- (A) $3x^3 - |x|$ (B) $|3x^2 - 1|$ (C) $3x^2|x| - 1$ (D) $3|x| - 1$ (E) $3x^2 - 1$

- _____ 29. If $e^{g(x)} = \frac{x^x}{x^2 - 1}$, then $g(x) =$
(A) $x \ln x - 2x$ (B) $\frac{\ln x}{2}$ (C) $(x-2)\ln x$ (D) $\frac{x \ln x}{\ln(x^2 - 1)}$ (E) $x \ln x - \ln(x^2 - 1)$

- _____ 30. $\frac{\ln(x^3 e^x)}{x} =$
(A) $\frac{3(\ln x + e^x)}{x}$ (B) $\ln(x^3 e^x - x)$ (C) $\ln x^2 + 1$ (D) $\frac{3 \ln x + x}{x}$ (E) $\frac{3 \ln x}{x}$

- _____ 31. If $f(g(x)) = \sec(x^3 + 4)$, $f(x) = \sec x^3$, and $g(x)$ is not an integer multiple of $\frac{\pi}{2}$, then
 $g(x) =$

- (A) $\sqrt[3]{x+4}$ (B) $\sqrt[3]{x-4}$ (C) $\sqrt[3]{x^3+4}$ (D) $\sqrt[3]{x}-4$ (E) $\sqrt[3]{x}+4$

- _____ 32. If $f(x) = \log_b x$, then $f(bx) =$
(A) $bf(x)$ (B) $f(b)f(x)$ (C) $1+f(x)$ (D) $xf(b)$ (E) $f(x)$

_____ 33. Which of the following statements is true?

- (A) $\log_{\frac{1}{2}} 2 < \log_{\frac{1}{\sqrt{2}}} 2$ (B) $\log_3(2+4) = \log_3 2 + \log_3 4$ (C) $\log 2 > \log 4$
(D) $\log_{\frac{1}{5}}(5\sqrt{5}) = \frac{2}{3}$ (E) $\log_{\frac{1}{2}} 2 - \log_{\frac{1}{2}} 4 = \log_{\frac{1}{2}} 2$